



碧云天生物技术/Beyotime Biotechnology
 订货热线: 400-1683301或800-8283301
 订货e-mail: order@beyotime.com
 技术咨询: info@beyotime.com
 网址: http://www.beyotime.com

E-Cadherin Mouse Monoclonal Antibody

产品编号	产品名称	包装
AF0138	E-Cadherin Mouse Monoclonal Antibody	50μl

产品简介:

来源	用途	交叉反应性	分子量
Mouse	WB, ICC, IHC, FC, IF	H, M	130KDa

WB, Western blot; IP, Immunoprecipitation; IF, Immunofluorescence; IHC, Immunohistochemistry; ICC, Immunocytochemistry; FC, Flow Cytometry; ELISA, Enzyme-linked Immunosorbent Assay; ChIP, Chromatin Immunoprecipitation Assay.

H, Human; M, Mouse; R, Rat; C, Chicken; Cw, Cow; Dg, Dog; Gp, Guinea pig; Hm, Hamster; Hr, Horse; Mk, Monkey; Pg, Pig; Rb, Rabbit; S, Sheep; Z, Zebrafish; All, all species expected.

- 配套提供了Western一抗稀释液, 可以用于Western检测或其它适当用途时的一抗稀释。
- 建议抗体使用时的稀释比例如下(实际使用时需根据抗原水平的高低作适当调整):

WB	IP	IF	IHC	ICC	FC	ELISA	ChIP
1:500-1:1000	-	1:200	1:200	1:200	1:50-1:100	-	-

- 抗体详细信息如下:

About this Antibody	
Name	E-Cadherin Mouse Monoclonal Antibody
Category	Monoclonal antibody(mAb); Primary antibody
Isotype	IgG1
Purification	ProA affinity purified
About the Immunogen	
Immunogen	This antibody is produced by immunizing mice with recombinant protein to a region of E-cadherin.
Gene ID	999
SwissProt	P12830
Synonyms	CADH1; cadherin 1; type 1; E-cadherin (epithelial) (ECAD); cadherin-1; CAM 120/80; E-cadherin; epithelial-cadherin; uvomorulin
Category	Adhesion/ECM
Background	E-cadherin (epithelial) is the most well-studied member of the cadherin family. It consists of 5 cadherin repeats (EC1 ~ EC5) in the extracellular domain, one transmembrane domain, and an intracellular domain that binds p120-catenin and beta-catenin. The intracellular domain contains a highly-phosphorylated region vital to beta-catenin binding and, therefore, to E-cadherin function. Loss of E-cadherin function or expression has been implicated in cancer progression and metastasis. E-cadherin downregulation decreases the strength of cellular adhesion within a tissue, resulting in an increase in cellular motility. This in turn may allow cancer cells to cross the basement membrane and invade surrounding tissues. E-cadherin is also used by pathologists to diagnose different kinds of breast cancer.

包装清单:

产品编号	产品名称	包装
AF0138	E-Cadherin Mouse Monoclonal Antibody	50μl
AZ050	Western一抗稀释液	50ml
—	说明书	1份

保存条件:

E-Cadherin Mouse Monoclonal Antibody -20°C保存, Western一抗稀释液-20°C或4°C保存, 一年有效。Western一抗稀释液优先推荐4°C保存, 长期不使用可以考虑-20°C保存, 但冻融可能会导致出现轻微的浑浊和少量不溶物。

注意事项:

- 如果本抗体用于Western blot (WB)、免疫荧光(IF)、免疫细胞化学(ICC)等实验, 请注意回收使用过的稀释抗体。回收的抗体通常至少可以重复使用5-10次。稀释后的抗体, 包括已经使用过的稀释抗体, 请4°C保存。
- 回收后重复使用的抗体, 使用方法同新鲜稀释的抗体。如果在重复使用过程中发现抗体出现轻微混浊现象, 可以10,000g离心1-3分钟, 取上清用于后续检测。如果回收的抗体出现明显的絮状物或长霉长菌等情况, 则可以考虑废弃该抗体。
- 提供的Western一抗稀释液也可以用于免疫荧光(IF)、免疫组化(IHC)、免疫细胞化学(ICC)等适当用途。如果希望获得最佳的检测效果, 请考虑使用上述检测专用的一抗稀释液。
- 本产品仅限于专业人员的科学研究用, 不得用于临床诊断或治疗, 不得用于食品或药品, 不得存放于普通住宅内。
- 为了您的安全和健康, 请穿实验服并戴一次性手套操作。

使用说明:

请根据抗体的实际用途选择相应的使用方法。

1. Western检测:

- 按照推荐的稀释比例用碧云天提供的Western一抗稀释液稀释抗体。
- 把经过封闭的蛋白膜与稀释好的一抗4°C缓慢摇动过夜或室温缓慢摇动2小时, 确保稀释的抗体至少能在摇动的瞬间覆盖蛋白膜。
- 回收稀释的一抗, 4°C保存以备下次继续使用。
- 按照Western的实验步骤进行后续的洗涤、二抗孵育、洗涤和检测等操作。具体操作可以参考如下网页:
<http://www.beyotime.com/support/western.htm>

2. 免疫染色:

可以使用碧云天生产的免疫染色一抗稀释液(P0103)稀释抗体, 使用后注意回收稀释好的一抗, 具体操作可以参考如下网页: <http://www.beyotime.com/support/immunol-staining.htm>

3. 其它实验操作请自行参考适当的protocol进行。

4. 代表性图片:




Fig. 1. Western blot analysis of E-cadherin on different cell lysates using anti- E-cadherin antibody at 1/1000 dilution. Positive control: Lane 1: A431 Lane 2: SW480 Lane 3: MCF-7 Lane 4: HeLa.




Fig. 2. Immunocytochemistry staining E-Cadherin in A431 cells (red). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X-100/PBS.




Fig. 3. Immunohistochemical analysis of paraffin-embedded human liver carcinoma tissue using E-Cadherin antibody. Counter stained with hematoxylin.




Fig. 4. Flow cytometric analysis of HeLa cells with E-Cadherin antibody at 1/50 dilution (blue) compared with an unlabelled control (cells without incubation with primary antibody; red). Goat anti mouse IgG

(FITC) was used as the secondary antibody.

相关产品:

产品编号	产品名称	包装
P0006	Bradford蛋白浓度测定试剂盒	1000次
P0010	BCA蛋白浓度测定试剂盒(增强型)	500次
P0012	BCA蛋白浓度测定试剂盒	500次
P0012A	SDS-PAGE凝胶配制试剂盒	1盒
P0012AC	SDS-PAGE凝胶快速配制试剂盒	1盒
P0013	Western及IP细胞裂解液	100ml
P0013B	RIPA裂解液(强)	100ml
P0014B	SDS-PAGE电泳液	10×1L
P0015	SDS-PAGE蛋白上样缓冲液(5X)	2ml
P0018	BeyoECL Plus(超敏ECL化学发光试剂盒)	共100ml
P0018A	BeyoECL Star(特超敏ECL化学发光试剂盒)	共100ml
P0020	显影定影试剂盒	各1升
P0021B	Western转膜液	10×1L
P0023A	Western一抗稀释液	100ml
P0023B	Western封闭液	100ml
P0023C6	Western洗涤液(10X)	10×100ml
P0023D	Western二抗稀释液	100ml
P0025	Western一抗二抗去除液	250ml
P0252	QuickBlock™ Western封闭液	100ml
P0256	QuickBlock™ Western一抗稀释液	100ml
P0258	QuickBlock™ Western二抗稀释液	100ml
P0066	预染蛋白质分子量标准	200μl
P0068	彩色预染蛋白质分子量标准	200μl
P0071	BeyoColor™彩色预染蛋白质分子量标准	200μl
P0098	免疫染色固定液	100ml
P0102	免疫染色封闭液	100ml
P0103	免疫染色一抗稀释液	100ml
P0106L	免疫染色洗涤液(10X)	10×100ml
P0108	免疫荧光染色二抗稀释液	100ml
P0110	免疫染色(非荧光)二抗稀释液	100ml
P0260	QuickBlock™免疫染色封闭液	100ml
P0262	QuickBlock™免疫染色一抗稀释液	100ml
P0265	QuickBlock™免疫荧光染色二抗稀释液	100ml
P0267	QuickBlock™免疫组化染色二抗稀释液	100ml
P0126	抗荧光淬灭封片液	5ml
FFP24	PVDF膜(进口分装, 6.6×8.5cm, 0.2μm)	20张/包装
FFP32	PVDF膜(进口原装, 6.6×8.5cm, 0.45μm)	20张/包装
FFP51	转印滤纸(7.5×10cm)	100片/包装
FF057	X-OMAT BT胶片(原柯达, 5×7英寸)	100张/盒
FFC58	压片暗盒(5×7英寸)	1个

使用本产品的文献:

1. Chi Shu, Haoran Zha, Haixia Long, Xinxin Wang, Fei Yang, Jianbao Gao, Chunyan Hu, Li Zhou, Bo Guo, Bo Zhu . C3a-C3aR signaling promotes breast cancer lung metastasis via modulating carcinoma associated fibroblasts J EXP CLIN CANC RES. 2020 Jan 13;39(1):11.
2. Huang C, Tao L, Wang XL, Pang Z . Berberine reversed the epithelial-mesenchymal transition of normal colonic epithelial cells induced by SW480 cells through regulating the important components in the TGF-β pathway. J Cell Physiol. 2018 Dec 10.
3. Dong Yuan, Yiyu Chen, Zhu Yang, Gang Li, Mingjun Wu, Jinyue Jiang, Dan Li, Qiubo Yu . SPOP attenuates migration and invasion of choriocarcinoma cells by promoting DHX9 degradation Am J Cancer Res. 2020 Aug 1;10(8):2428-2445.
4. Wu J, Gao F, Xu T, Deng X, Wang C, Yang X, Hu Z, Long Y, He X, Liang G, Ren D, Dai T . miR-503 suppresses the proliferation and metastasis of esophageal squamous cell carcinoma by triggering autophagy via PKA/mTOR signaling. Int J Oncol. 2018 Mar 16.

5. Zhang J, Chen D, Liang S, Wang J, Liu C, Nie C, Shan Z, Wang L, Fan Q, Wang F . miR-106b promotes cell invasion and metastasis via PTEN mediated EMT in ESCC. *Oncol Lett*. 2018 Apr;15(4):4619-4626.
6. Liu Z, Wu Y, Tao Z, Ma L . E3 ubiquitin ligase Hakai regulates cell growth and invasion, and increases the chemosensitivity to cisplatin in non-small-cell lung cancer cells. *Int J Mol Med*. 2018 Aug;42(2):1145-1151.
7. Feng Y, Guo X, Huang X, Wu M, Li X, Wu S, Luo X . Metformin reverses stem cell-like HepG2 sphere formation and resistance to sorafenib by attenuating epithelial-mesenchymal transformation. *Mol Med Rep*. 2018 Oct;18(4):3866-3872.
8. Yang W, Gong X, Wang X, Huang C . A mediator of phosphorylated Smad2/3, evodiamine, in the reversion of TAF-induced EMT in normal colonic epithelial cells. *INVEST NEW DRUG*. 2018 Nov 29.
9. Wang XL, Huang C . Difference of TGF- β /Smads signaling pathway in epithelial-mesenchymal transition of normal colonic epithelial cells induced by tumor-associated fibroblasts and colon cancer cells. *Mol Biol Rep*. 2019 Jun 46(3):2749-2759.
10. Zhang T, Liu C, Yu Y, Geng J, Meng Q, Xu S, Zhou F, Chen Y, Jin S, Shen J, Pan B, Meng F, Liu F . TBL1XR1 is involved in c-Met-mediated tumorigenesis of human nonsmall cell lung cancer. *Cancer Gene Ther*. 2019 Jun 27
11. Lu Y, Liang M, Zhang Q, Liu Z, Song Y, Lai L, Li Z . Mutations of GADD45G in rabbits cause cleft lip by the disorder of proliferation, apoptosis and epithelial-mesenchymal transition (EMT). *BBA-MOL BASIS DIS*. 2019 Sep 1 1865(9):2356-2367.
12. Wang K, Zou P, Zhu X, Zhang T . Ziyuglycoside II suppresses the aggressive phenotype of triple negative breast cancer cells through regulating Src/EGFR-dependent ITGB4/FAK signaling. *Toxicol In Vitro*. 2019 Dec 61:104653.
13. Pan H, Xue W, Zhao W, Schachner M. . Expression and function of chondroitin 4-sulfate and chondroitin 6-sulfate in human glioma. *FASEB J*. 2020 Feb 34(2):2853-2868.
14. Tiewa Zhang, Cheng Liu, Yan Yu, Jianxiong Geng, Qingwei Meng, Shanqi Xu, Fengrui Zhou, Yingying Chen, Shi Jin, Jing Shen, Bo Pan, Fanling Meng, Fang Liu . TBL1XR1 is involved in c-Met-mediated tumorigenesis of human nonsmall cell lung cancer *Cancer Gene Ther*. 2020 Apr;27(3-4):136-146.
15. Shaohan Zou, Ruirui Dong, Ping Zou, Xina Meng, Ting Zhang, Liang Luo, Na Li, Yao Wang, Jing Wang, Tiejun Wang, Yan Zhang, Minjian Chen, Conghua Zhou, Fei Han . ERp29 affects the migratory and invasive ability of human extravillous trophoblast HTR-8/SVneo cells via modulating the epithelial-mesenchymal transition *J BIOCHEM MOL TOXIC*. 2020 Apr;34(4):e22454.
16. Chao Huang, Ke Ming Xiang, Bing Jun Liang, Wei Xuan Huang, Fan Jun Zhang, Yu Wan Shao, Xiu Lian Wang, Hao Sheng Liu, Wei Zeng Shen . Combination of Evodiamine with Berberine Reveals a Regulatory Effect on the Phenotypic Transition of Colon Epithelial Cells Induced by CCD-18Co *J TRADIT CHIN MED*. 2020 Sep 30;35(3):195-206.
17. Sijia Lei, Bin Zhang, Luyuan Huang, Ziyu Zheng, Shaohan Xie, Lianghua Shen, Mason Breitzig, Alexander Czachor, Hongtao Liu, Huiru Luo, Yanxia Chen, Kangshou Liu, Hanxiao Sun, Qing Zheng, Qiang Li, Feng Wang . SRSF1 promotes the inclusion of exon 3 of SRA1 and the invasion of hepatocellular carcinoma cells by interacting with exon 3 of SRA1pre-mRNA *Cell Death Discov*. 2021 May 19;7(1):117.
18. Chengwu Tang, Wenming Feng, Ying Bao, Huimin Du . Long non-coding RNA TINCR promotes hepatocellular carcinoma proliferation and invasion via STAT3 signaling by direct interacting with T-cell protein tyrosine phosphatase (TCPTP) *Bioengineered*. 2021 Dec;12(1):2119-2131.
19. Lili Zhu, Xiaobei Yang, Jingyu Feng, Jian Mao, Qidong Zhang, Mengru He, Yang Mi, Yingwu Mei, Ge Jin, Haifeng Zhang . CYP2E1 plays a suppressive role in hepatocellular carcinoma by regulating Wnt/Dvl2/ β -catenin signaling *J Transl Med*. 2022 May 4;20(1):194.

Version 2024.03.12