



碧云天生物技术/Beyotime Biotechnology  
 订货热线: 400-168-3301或800-8283301  
 订货e-mail: order@beyotime.com  
 技术咨询: info@beyotime.com  
 网址: http://www.beyotime.com

## Ubiquitin Rabbit Polyclonal Antibody

| 产品编号   | 产品名称                                 | 包装   |
|--------|--------------------------------------|------|
| AF0306 | Ubiquitin Rabbit Polyclonal Antibody | 50μl |

### 产品简介:

| 来源     | 用途               | 交叉反应性   | 分子量   |
|--------|------------------|---------|-------|
| Rabbit | WB, ICC, IHC, IF | H, M, R | 10KDa |

WB, Western blot; IP, Immunoprecipitation; IF, Immunofluorescence; IHC, Immunohistochemistry; ICC, Immunocytochemistry; FC, Flow Cytometry; ELISA, Enzyme-linked Immunosorbent Assay; ChIP, Chromatin Immunoprecipitation Assay.

H, Human; M, Mouse; R, Rat; C, Chicken; Cw, Cow; Dg, Dog; Gp, Guinea pig; Hm, Hamster; Hr, Horse; Mk, Monkey; Pg, Pig; Rb, Rabbit; S, Sheep; Z, Zebrafish; All, all species expected.

- 配套提供了Western一抗稀释液, 可以用于Western检测或其它适当用途时的一抗稀释。
- 建议抗体使用时的稀释比例如下(实际使用时需根据抗原水平的高低作适当调整):

| WB           | IP | IF    | IHC   | ICC   | FC | ELISA | ChIP |
|--------------|----|-------|-------|-------|----|-------|------|
| 1:500-1:1000 | -  | 1:200 | 1:200 | 1:200 | -  | -     | -    |

- 抗体详细信息如下:

| About this Antibody |  |
|---------------------|--|
| Name                | Ubiquitin Rabbit Polyclonal Antibody   |
| Category            | Polyclonal antibody(pAb); Primary antibody   |
| Isotype             | IgG  |
| Purification        | Peptide affinity purified  |
| About the Immunogen |  |
| Immunogen           | This antibody is produced by immunizing rabbits with a synthetic peptide (KLH-coupled) corresponding to Ubiquitin.   |
| Gene ID             | 7314   |
| SwissProt           | P62988   |
| Synonyms            | 40S ribosomal protein S27a; CEP80; Ribosomal protein S27a; RPS27A; UBA52; UBA80; UBB; UBC; UBCEP1; UBCEP80; UBIQ; ubiquitin and ribosomal protein S27a; ubiquitin carboxyl extension protein 80; ubiquitin-CEP80   |
| Category            | Ubiquitin and Ubiquitin-like proteins  |
| Background          | Ubiquitin is a conserved polypeptide unit that plays an important role in the ubiquitin-proteasome pathway. Ubiquitin exists either covalently attached to another protein, or free (unanchored). When covalently bound, it is conjugated to target proteins via an isopeptide bond either as a monomer (monoubiquitin), a polymer linked via different Lys residues of the ubiquitin (polyubiquitin chains) or a linear polymer linked via the initiator Met of the ubiquitin (linear polyubiquitin chains). Polyubiquitin chains, when attached to a target protein, have different functions depending on the Lys residue of the ubiquitin that is linked: Lys-6-linked may be involved in DNA repair; Lys-11-linked is involved in ERAD (endoplasmic reticulum-associated degradation) and in cell-cycle regulation; Lys-29-linked is involved in lysosomal degradation; Lys-33-linked is involved in kinase modification; Lys-48-linked is involved in protein degradation via the proteasome; Lys-63-linked is involved in endocytosis, DNA-damage responses as well as in signaling processes leading to activation of the transcription factor NF-kappa-B. |

### 包装清单:

| 产品编号   | 产品名称                                 | 包装   |
|--------|--------------------------------------|------|
| AF0306 | Ubiquitin Rabbit Polyclonal Antibody | 50μl |
| AZ050  | Western一抗稀释液                         | 50ml |
| -      | 说明书                                  | 1份   |

## 保存条件:

Ubiquitin Rabbit Polyclonal Antibody -20°C保存, Western一抗稀释液-20°C或4°C保存, 一年有效。Western一抗稀释液优先推荐4°C保存, 长期不使用可以考虑-20°C保存, 但冻融可能会导致出现轻微的浑浊和少量不溶物。

## 注意事项:

- 如果本抗体用于Western blot (WB)、免疫荧光(IF)、免疫细胞化学(ICC)等实验, 请注意回收使用过的稀释抗体。回收的抗体通常至少可以重复使用5-10次。稀释后的抗体, 包括已经使用过的稀释抗体, 请4°C保存。
- 回收后重复使用的抗体, 使用方法同新鲜稀释的抗体。如果在重复使用过程中发现抗体出现轻微混浊现象, 可以10,000g离心1-3分钟, 取上清用于后续检测。如果回收的抗体出现明显的絮状物或长霉长菌等情况, 则可以考虑废弃该抗体。
- 提供的Western一抗稀释液也可以用于免疫荧光(IF)、免疫组化(IHC)、免疫细胞化学(ICC)等适当用途。如果希望获得最佳的检测效果, 请考虑使用上述检测专用的一抗稀释液。
- 本产品仅限于专业人员的科学研究用, 不得用于临床诊断或治疗, 不得用于食品或药品, 不得存放于普通住宅内。
- 为了您的安全和健康, 请穿实验服并戴一次性手套操作。

## 使用说明:

请根据抗体的实际用途选择相应的使用方法。

### 1. Western检测:

- 按照推荐的稀释比例用碧云天提供的Western一抗稀释液稀释抗体。
- 把经过封闭的蛋白膜与稀释好的一抗4°C缓慢摇动过夜或室温缓慢摇动2小时, 确保稀释的抗体至少能在摇动的瞬间覆盖蛋白膜。
- 回收稀释的一抗, 4°C保存以备下次继续使用。
- 按照Western的实验步骤进行后续的洗涤、二抗孵育、洗涤和检测等操作。具体操作可以参考如下网页:  
<http://www.beyotime.com/support/western.htm>

### 2. 免疫染色:

可以使用碧云天生产的免疫染色一抗稀释液(P0103)稀释抗体, 使用后注意回收稀释好的一抗, 具体操作可以参考如下网页: <http://www.beyotime.com/support/immunol-staining.htm>

### 3. 其它实验操作请自行参考适当的protocol进行。

### 4. 代表性图片:




Fig. 1. Western blot analysis of Ubiquitin on different cell lysates using anti-Ubiquitin antibody at 1/1000 dilution.

Positive control: Lane 1: 293T Lane 2: HepG2 Lane 3: HeLa Lane 4: MCF-2.




Fig. 2. Immunocytochemistry staining Ubiquitin in MCF-7 cells (green). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X-100/PBS.




Fig. 3. Immunohistochemical analysis of paraffin-embedded mouse spleen tissue using anti-Ubiquitin antibody. Counter stained with hematoxylin.

## 相关产品:

| 产品编号  | 产品名称              | 包装    |
|-------|-------------------|-------|
| P0006 | Bradford蛋白浓度测定试剂盒 | 1000次 |
| P0010 | BCA蛋白浓度测定试剂盒(增强型) | 500次  |

|         |                                |          |
|---------|--------------------------------|----------|
| P0012   | BCA蛋白浓度测定试剂盒                   | 500次     |
| P0012A  | SDS-PAGE凝胶配制试剂盒                | 1盒       |
| P0012AC | SDS-PAGE凝胶快速配制试剂盒              | 1盒       |
| P0013   | Western及IP细胞裂解液                | 100ml    |
| P0013B  | RIPA裂解液(强)                     | 100ml    |
| P0014B  | SDS-PAGE电泳液                    | 10×1L    |
| P0015   | SDS-PAGE蛋白上样缓冲液(5X)            | 2ml      |
| P0018   | BeyoECL Plus(超敏ECL化学发光试剂盒)     | 共100ml   |
| P0018A  | BeyoECL Star(特超敏ECL化学发光试剂盒)    | 共100ml   |
| P0020   | 显影定影试剂盒                        | 各1升      |
| P0021B  | Western转膜液                     | 10×1L    |
| P0023A  | Western一抗稀释液                   | 100ml    |
| P0023B  | Western封闭液                     | 100ml    |
| P0023C6 | Western洗涤液(10X)                | 10×100ml |
| P0023D  | Western二抗稀释液                   | 100ml    |
| P0025   | Western一抗二抗去除液                 | 250ml    |
| P0252   | QuickBlock™ Western封闭液         | 100ml    |
| P0256   | QuickBlock™ Western一抗稀释液       | 100ml    |
| P0258   | QuickBlock™ Western二抗稀释液       | 100ml    |
| P0066   | 预染蛋白质分子量标准                     | 200μl    |
| P0068   | 彩色预染蛋白质分子量标准                   | 200μl    |
| P0071   | BeyoColor™彩色预染蛋白质分子量标准         | 200μl    |
| P0098   | 免疫染色固定液                        | 100ml    |
| P0102   | 免疫染色封闭液                        | 100ml    |
| P0103   | 免疫染色一抗稀释液                      | 100ml    |
| P0106L  | 免疫染色洗涤液(10X)                   | 10×100ml |
| P0108   | 免疫荧光染色二抗稀释液                    | 100ml    |
| P0110   | 免疫染色(非荧光)二抗稀释液                 | 100ml    |
| P0260   | QuickBlock™免疫染色封闭液             | 100ml    |
| P0262   | QuickBlock™免疫染色一抗稀释液           | 100ml    |
| P0265   | QuickBlock™免疫荧光染色二抗稀释液         | 100ml    |
| P0267   | QuickBlock™免疫组化染色二抗稀释液         | 100ml    |
| P0126   | 抗荧光淬灭封片液                       | 5ml      |
| FFP24   | PVDF膜(进口分装, 6.6×8.5cm, 0.2μm)  | 20张/包装   |
| FFP32   | PVDF膜(进口原装, 6.6×8.5cm, 0.45μm) | 20张/包装   |
| FFP51   | 转印滤纸(7.5×10cm)                 | 100片/包装  |
| FF057   | X-OMAT BT胶片(原柯达, 5×7英寸)        | 100张/盒   |
| FFC58   | 压片暗盒(5×7英寸)                    | 1个       |

Version 2016.04.26