



Regulation of cellular glutathione peroxidase by different forms and concentrations of selenium in primary cultured bovine hepatocytes

Xianshi Wu^{a,b}, Kehe Huang^{a,*}, Chengwu Wei^a, Fu Chen^a, Cuiling Pan^a

^a*Institute of Nutritional and Metabolic Disorders in Domestic Animals and Fowls, Nanjing Agricultural University, Nanjing 210095, China*

^b*College of Animal Science and Technology, Guangxi University, Nanning 530004, China*

Received 22 May 2008; received in revised form 7 December 2008; accepted 10 December 2008

Abstract

The expression and activity of cellular glutathione peroxidase (GPx1) are regulated by selenium (Se). Generally speaking, organic forms of Se have less toxicity and greater bioavailability compared with inorganic forms. In this study, the effects of different forms and concentrations of Se on the regulation of mRNA level and activity of GPx1 in bovine hepatocytes were evaluated, and the optimal doses of different forms of Se that supported the full expression of GPx1 were determined. Primary cultured bovine hepatocyte monolayers derived from neonatal male Holstein calves (aged 1–2 days) were incubated for 24 h with 0 (control), 0.5, 1, 1.5, 2, 3, 4 or 5 $\mu\text{mol/L}$ of Se from DL-selenomethionine (Se-Met), sodium selenite (Na_2SeO_3) or Kappa-selenocarrageenan (Se-Car). Compared with controls, a significantly lower level of release of lactic dehydrogenase (LDH) was observed at 0.5–5 $\mu\text{mol/L}$ of Se-Met, 0.5–1 $\mu\text{mol/L}$ of Na_2SeO_3 and 0.5 $\mu\text{mol/L}$ of Se-Car, but significantly higher LDH release was observed at 2–5 $\mu\text{mol/L}$ of Na_2SeO_3 and 3–5 $\mu\text{mol/L}$ of Se-Car, and the response occurred in a dose-dependent manner. The intracellular content of reduced glutathione in all hepatocytes treated with Se was significantly lower than that of controls. Significant increases in GPx1 mRNA were obtained in all hepatocytes treated with Se, with maximal effects at 3 $\mu\text{mol/L}$ of Se-Met, 1.5 $\mu\text{mol/L}$ of Na_2SeO_3 and 2 $\mu\text{mol/L}$ of Se-Car, respectively. Furthermore, 3 $\mu\text{mol/L}$ of Se from Se-Met resulted in peak levels of GPx1 mRNA. After reaching a maximal level, higher Se supplementation led to a reduction of GPx1 mRNA. The activity of GPx1 showed similar patterns but of lower magnitude. We conclude that (a) the regulation of mRNA level and activity of GPx1 in primary cultured bovine hepatocytes by different forms of Se varies and (b) the optimal doses of Se to support the full expression of GPx1 in bovine hepatocytes when supplied as Se-Met, Na_2SeO_3 and Se-Car are 3, 1.5 and 2 $\mu\text{mol/L}$, respectively.

© 2009 Published by Elsevier Inc.

Keywords: Se; GPx1; Bovine hepatocytes; Gene expression; Bioavailability

1. Introduction

Selenium (Se) is an essential trace element for animals and humans. Previous data suggest that Se may enhance immune function [1], reproductive performance [2,3] and resistance to oxidative stress [4]. Supplementation of Se can be protective, to some extent, against certain types of cancer [5–8]. Deficiency of Se can be associated with enhanced severity of infectious diseases [1,9] and is linked with numerous diseases such as Keshan disease and Kaschin–Beck disease in humans [10,11], “white muscle disease” in calves and lambs [12] and exudative diathesis in chicks [13].

Cellular glutathione peroxidase (GPx1, E.C. 1.11.1.9), a homotetramer of 84 kDa, was the first Se-dependent enzyme to be identified. The mRNA sequence of GPx1 contains a UGA codon that encodes a selenocysteine (Sec) residue, and this is normally used as a translation termination codon. The Sec residue is located at the active center of GPx1 [14,15]. Supplementation with Se may increase the stability of cytoplasmic GPx1 mRNA [16]. In contrast, Se deficiency decreases the efficiency of incorporation of Sec at the UGA codon site during mRNA translation and results in decay of cytoplasmic GPx1 mRNA, which is mediated by the nonsense codon [17].

Reactive oxygen species (ROS) are generated primarily by electron leakage from mitochondrial electron carriers and enzymes during oxidative phosphorylation in aerobic mammalian cells. If ROS are not removed in a timely

* Corresponding author. Tel.: +86 25 84395507; fax: +86 25 84398669.
E-mail address: khhuang@njau.edu.cn (K. Huang).

fashion by an antioxidant enzyme system and overstep physiological concentrations, the cells may encounter oxidative stress, which causes destruction of lipids, proteins and DNA. It is known that GPx, along with superoxide dismutase and catalase, forms the main antioxidant enzyme system in mammals. GPx1 can detoxify hydrogen peroxide to water and a range of organic peroxides to the corresponding alcohols using reduced glutathione (GSH) as the hydrogen donor [15]. Therefore, GPx1 acts as one of the cellular protectors against oxidative damage.

Previous studies have suggested that organic forms of Se [e.g., Se-enriched yeast, which contains selenomethionine (Se-Met)] are less toxic than inorganic forms of Se such as sodium selenite [18–20]. Organic forms of Se have excellent bioavailability and are appropriate for nutritional supplementation [21,22]. Because the expression and activity of GPx1 are highly correlated with the Se status of the animal or cell [23,24], GPx1 has been used as a biochemical indicator of Se status and requirements [25]. Lei et al. [26] reported that Se supplementation at 0.2 mg Se/kg of diet was required to support the full expression of GPx1 in the livers of young pigs. In addition, Sunde et al. [27] showed that maximal activity of GPx1 in the livers of pregnant and lactating rats required 0.075 and 0.1 mg Se/kg of diet, respectively.

The diets of dairy cattle are routinely supplemented with Se. Various inorganic and organic forms of Se are used in many countries. There are no established criteria for the levels of Se that are released from inorganic or organic forms in the rations fed to Chinese dairy cattle. It is known that the mammalian liver is the principal organ involved in storage of Se and that it can synthesize GPx1, GPx4, thioredoxin reductase and deiodinase. However, little is known about the regulation of mRNA level and activity of GPx1 in bovine hepatocytes. The purposes of the present study were to evaluate the effects of different forms and concentrations of Se on the regulation of mRNA level and activity of GPx1 in primary cultured bovine hepatocytes and to determine the optimal doses of different forms of Se for support of the full expression of GPx1 mRNA *in vitro*.

2. Materials and methods

2.1. Preparation of bovine hepatocytes

Bovine livers were obtained from neonatal male Holstein calves (aged 1–2 days), which were used to produce serum and slaughtered at the local slaughterhouse. The carcass of each calf was skinned and the abdominal cavity was opened by butchers. The caudate process of the liver was excised by a single transverse cut with sterile instruments and placed on a polypropylene tray with the hepatic vein uppermost. There were two large (4–6 mm in diameter) hepatic vessels on the cut surface. Blood was removed from the organ by perfusion with calcium-free cold HEPES buffer (33 mM HEPES, 127.8 mM NaCl, 3.15 mM KCl, 0.7 mM Na₂HPO₄ and 0.6 mM EGTA, pH 7.65) through

the hepatic vessels. The liver lobe was then immersed in physiological saline (1–4°C) and transported to the laboratory. The interval from death of the calf to hepatocyte isolation was less than 1 h.

Hepatocyte isolation was performed by the collagenase perfusion method as previously described [28,29], with some modifications. The liver lobe was rewarmed by preperfusion for 10–12 min with the same calcium-free HEPES buffer (37–38°C) but without EGTA. The liver lobe was then perfused with 300 ml of 0.05% (w/v) collagenase type II (Invitrogen, USA) perfusion buffer (33 mM HEPES, 127.8 mM NaCl, 3.15 mM KCl, 0.7 mM Na₂HPO₄ and 3 mM CaCl₂, pH 7.65, 37–38°C) using a recirculating system at a flow rate of approximately 120 ml/min for 20–25 min. Digested liver tissue was then transferred to a sterile beaker and the liver capsule was removed. The liver tissue was disrupted gently and the cells were dispersed in 40 ml of serum-free DMEM/Ham F12 medium (Invitrogen) supplemented with 0.2% (w/v) bovine serum albumin (BSA) and antibiotics (100 IU/ml penicillin, 100 µg/ml streptomycin and 0.25 µg/ml amphotericin B) at 4°C. The cell suspension was filtered once through each mesh (grid size about 150 and 75 µm, respectively) and further diluted with 300 ml of the same serum-free medium. After sedimentation for 10 min at 4°C, the supernatant was discarded. The cells were pelleted by centrifugation (30×g and 50×g for 3 min each) and then resuspended and washed twice with the same serum-free medium at 4°C. Finally, the isolated cells were resuspended in 40 ml of DMEM/Ham F12 medium containing 15% fetal bovine serum (FBS), 24 mM NaHCO₃, 2 mM L-glutamine, 0.2% (w/v) BSA, 100 nM dexamethasone, 5 mg/L transferrin, 1 µmol/L insulin and antibiotics. The viability of isolated hepatocytes was 83.75±1.78% (mean±S.D., n=3) by 0.4% trypan blue dye exclusion, and the cell yield from each liver lobe preparation was 3.34±0.71×10⁸ hepatocytes (mean±S.D., n=3). Evaluation of the cells by light microscopy indicated that about 94% of the collected cells were hepatocytes.

2.2. Hepatocyte cultures and treatments

Hepatocytes were seeded into six-well plates precoated with cattle tail tendon collagen at a density of 1.2×10⁶ viable cells per well in 2 ml of DMEM/Ham F12 medium. Cells were cultured at 37°C and 5% CO₂ in a humidified atmosphere. After an attachment period of 4 h, cell monolayers were washed twice with Hanks' balanced salt solution, and 3 ml of fresh DMEM/Ham F12 medium containing 10% FBS was added to each well. Following culture for 24 h, the culture medium was removed and the cell monolayers were washed three times with Hanks' balanced salt solution. Then, hepatocytes were grown in 3 ml of fresh serum-free DMEM/Ham F12 medium and treated with 0 (control), 0.5, 1, 1.5, 2, 3, 4 or 5 µmol/L of Se as DL-Se-Met (Sigma, USA), sodium selenite (Na₂SeO₃, Sigma) or Kappa-selenocarrageenan (Se-Car, First Institute of

Oceanography, China). After 24 h of incubation, the culture medium was collected into 2-ml Eppendorf tubes for assay of lactic dehydrogenase (LDH) and cells were harvested for analysis of mRNA and enzyme activity. Each treatment was done in six wells (three wells for mRNA analysis, the others for measurement of GPx1 activity and GSH concentration) in three separate experiments.

2.3. Preparation of hepatocyte lysates

After 24 h incubation with different forms and concentrations of Se, the cell monolayers were washed three times with ice-cold $\text{Ca}^{2+}/\text{Mg}^{2+}$ -free phosphate-buffered saline (PBS) and harvested by scraping into 1 ml of cold Tris buffer (20 mM Tris-HCl, pH 7.5, 2 mM EDTA and 0.1% peroxide-free Triton X-100). Hepatocyte lysates were then prepared by ultrasonication for 30 s in icy water and centrifuged at $12,000\times g$ for 15 min at 4°C . The supernatants were aliquoted and stored at -20°C for subsequent analysis.

2.4. Analytical methods

2.4.1. Determination of protein concentration of hepatocyte lysates

Total protein concentration was determined using a Bradford Protein Assay Kit (Beyotime Institute of Biotechnology, China). The enzyme activity results were corrected for total protein concentration.

2.4.2. Determination of LDH activity in the culture medium

Hepatocyte toxicity was analyzed by measuring the activity of LDH in the incubation medium. The culture medium was collected into 2-ml Eppendorf tubes at the end of incubation and centrifuged at $12,000\times g$ for 15 min at 4°C . The supernatants were stored at -20°C until analysis. The measurement of LDH activity in the culture medium was performed as previously described [30]. One unit of enzyme activity was defined as 1 μmol of reduced nicotinamide adenine dinucleotide oxidized per minute. The activity of LDH in the culture medium was expressed as units per liter. All samples were measured in duplicate.

2.4.3. Measurement of GSH concentration in hepatocyte cytosol

The concentration of GSH in hepatocyte cytosol was measured according to the method described by Rahman et al. [31]. The method is based upon a reaction of 5,5'-dithiobis(2-nitrobenzoic acid) (Ellman's reagent) and GSH. The chromophoric product (5-thio-2-nitrobenzoic acid) possesses a molar absorption at a wavelength of 412 nm

that is proportional to the GSH concentration in the sample. The GSH concentrations were calculated from a standard curve prepared with pure GSH standards and were expressed as micromoles of GSH per gram of protein. All samples were assayed in duplicate.

2.4.4. Determination of GPx1 mRNA level by quantitative RT-PCR

After 24 h incubation with different forms and concentrations of Se, the medium was removed and the cell monolayers were washed three times with ice-cold $\text{Ca}^{2+}/\text{Mg}^{2+}$ -free PBS. Total RNA was isolated from the hepatocyte monolayers using the TRIZOL reagent (Invitrogen) according to the manufacturer's protocol. The dried RNA pellets were resuspended in 40 μl of diethyl-pyrocyanate-treated water. The concentration and purity of the total RNA were determined spectrophotometrically at 260/280 nm. The total RNA was immediately used or stored at -70°C prior to cDNA synthesis.

First-strand cDNA was synthesized from 2 μg of total RNA using Oligo dT primers and Superscript II reverse transcriptase (Invitrogen) according to the manufacturer's instructions. Synthesized cDNA was diluted 10 times with sterile water and stored at -20°C before use.

Primer Premier software (PREMIER Biosoft International, USA) was used to design specific primers for GPx1 and β -actin based on known bovine sequences (Table 1). General PCRs were first performed to confirm the specificity of the primers. The PCR products were electrophoresed on 2% agarose gels, extracted, cloned into the pMD18-T vector (TaKaRa, Dalian, China) and sequenced. The plasmids were serially diluted from 10^9 to 10^4 copies and used as templates in quantitative real-time PCRs to establish standard curves. Quantitative real-time PCR was performed on an ABI PRISM 7300 Detection System (Applied Biosystems, USA). Reactions were performed in a 25- μl reaction mixture containing 12.5 μl of $2\times$ SYBR Green I PCR Master Mix (TOYOBO, Japan), 10 μl of either diluted cDNA or plasmid standard, 1 μl of each primer (10 $\mu\text{mol}/\text{L}$) and 0.5 μl of PCR-grade water. The PCR procedure for GPx1 and β -actin consisted of a 95°C step for 2 min followed by 40 cycles consisting of 95°C for 15 s, 62°C for 15 s and 72°C for 30 s. The melting curve analysis showed only one peak for each PCR product. Electrophoresis was performed with the PCR products to verify primer specificity and the purity of the products. The calculation of the number of copies of each sample was performed from the respective standard curve using the

Table 1
Primers used for quantitative real-time PCR

Primers	Sequence (5'-3')	Target gene	GenBank accession no.	PCR fragment length (bp)
GPx1-F (forward)	ctt gct gct tgg cgg tca	GPx1	X13684	139
GPx1-R (reverse)	agg gga ggc tgg gat gga ta			
β -Actin-F (forward)	tgc ctt ccc aaa agc cac c	β -actin	BC102948	250
β -Actin-R (reverse)	acc tca acc cgc tcc caa			




Fig. 1. Effects of different forms and concentrations of Se on LDH release from hepatocytes. The primary cultured bovine hepatocyte monolayers were incubated with 0 (control), 0.5, 1, 1.5, 2, 3, 4 or 5 $\mu\text{mol/L}$ of Se as Se-Met, Na_2SeO_3 or Se-Car for 24 h. The activity of LDH in the culture medium was measured to evaluate cell membrane integrity. Bars represent mean \pm S.D. of triplicate culture media. Bars with “***” are statistically significantly different from control by one-way ANOVA followed by Tukey’s multiple comparison test ($*P<0.05$, $**P<0.01$, $***P<0.001$). Within the groups treated with different levels of Se-Met, Na_2SeO_3 or Se-Car, bars sharing a common letter are not significantly different ($P>0.05$). “x” and “xx” denote significant difference ($*P<0.05$, $^{xx}P<0.01$) between cells treated with Na_2SeO_3 and Se-Car.

7300 system software. The ratio of the level of GPx1 mRNA to that of the β -actin internal control was used for statistical comparison of the different treatments.

2.4.5. Determination of GPx1 activity in hepatocyte cytosol

The activity of GPx1 in hepatocyte cytosol was measured according to the method described by Lei et al. [32], with *tert*-butyl hydroperoxide (*t*-Bu-OOH) as the peroxide substrate. Briefly, 50 μl of hepatocyte lysate was transferred to a 3-ml quartz cuvette containing 1900 μl of the reaction mixture [50 mM Tris-HCl, pH 7.5, 2 mM EDTA, 0.1 mM reduced nicotinamide adenine dinucleotide phosphate (NADPH), 2 mM GSH, 1 mM NaN_3 and 0.9 U of GSH reductase (Sigma)]. The reaction mixture was preincubated for 3 min at 25°C, and the reaction was initiated by adding 50 μl of *t*-Bu-OOH (8 mM). The rate of oxidation of NADPH was monitored in a spectrophotometer at 340 nm for 5 min at 25°C. The nonenzymatic reaction rate was determined by substituting water (serving as the blank) for the hepatocyte lysate and recording the decrease in NADPH absorbance. One unit of enzyme activity was defined as 1 μmol of NADPH oxidized per minute under these conditions. The activity of GPx1 in hepatocyte cytosol was expressed as units per gram of protein. All samples were measured in duplicate.

2.5. Statistical analysis

Statistical analysis of LDH activity, GSH concentration, GPx1 activity and mRNA level was performed using the SPSS 11.5 for Windows statistical software package (SPSS Inc., USA). When a significant value ($P<0.05$) was obtained

by one-way analysis of variance (ANOVA), further analysis was done. All data showed a normal distribution and passed equal variance testing. Differences between means were assessed by the Tukey’s honestly significant difference test of post hoc multiple comparisons. Data were expressed as mean \pm S.D. Differences were considered as significant at $P<0.05$.

3. Results

3.1. LDH release into the culture medium

The effects of different forms and concentrations of Se on release of LDH into the culture medium are shown in Fig. 1. A significant reduction in LDH release was observed for primary cultured bovine hepatocytes incubated with Se-Met at doses of 0.5–5 $\mu\text{mol/L}$ for 24 h ($P<0.01$, vs. control). In the hepatocytes incubated with Na_2SeO_3 , significantly lower LDH release was observed at doses of 0.5 ($P<0.01$) and 1 ($P<0.05$) $\mu\text{mol/L}$ (vs. control) and significantly higher LDH release was observed at doses of 2, 3, 4 or 5 $\mu\text{mol/L}$ ($P<0.01$, vs. control); the response occurred in a dose-dependent manner. The release of LDH was significantly reduced by Se-Car at a dose of 0.5 $\mu\text{mol/L}$ ($P<0.01$), but not at doses of 1, 1.5 or 2 $\mu\text{mol/L}$ (vs. control). In contrast, significantly increased release of LDH was measured when hepatocytes were supplemented with 3, 4 or 5 $\mu\text{mol/L}$ of Se-Car ($P<0.01$, vs. control). Furthermore, the release of LDH induced by Na_2SeO_3 at doses of 3, 4 or 5 $\mu\text{mol/L}$ was significantly higher than that induced by Se-Car at equivalent doses. Cell detachments were not observed by light microscopy in the hepatocytes incubated with Na_2SeO_3 at a dose of 5 $\mu\text{mol/L}$.




Fig. 2. Effects of different forms and concentrations of Se on GSH concentration in hepatocyte cytosol. Hepatocyte cytosol was obtained from primary cultured bovine hepatocyte monolayers supplemented with 0 (control), 0.5, 1, 1.5, 2, 3, 4 or 5 µmol/L of Se as Se-Met, Na₂SeO₃ or Se-Car for 24 h. Total GSH concentration in hepatocyte cytosol was measured by a spectrophotometric method. Bars represent mean±S.D. of triplicate cultures. Bars with “****” are statistically significantly different from control by one-way ANOVA followed by Tukey’s multiple comparison test (***) ($P < 0.001$). Within the groups treated with different levels of Se-Met, Na₂SeO₃ or Se-Car, bars sharing a common letter are not significantly different ($P > 0.05$).




Fig. 3. Effects of different forms and concentrations of Se on GPx1 mRNA in bovine hepatocytes. The primary cultured bovine hepatocyte monolayers were treated with 0 (control), 0.5, 1, 1.5, 2, 3, 4 or 5 µmol/L of Se as Se-Met, Na₂SeO₃ or Se-Car for 24 h. GPx1 mRNA in bovine hepatocytes was measured by quantitative real-time RT-PCR, and the ratio of the level of GPx1 mRNA to that of the β-actin internal control was used for statistical comparison. Bars represent mean±S.D. of triplicate cultures. Bars with “****” are statistically significantly different from control by one-way ANOVA followed by Tukey’s multiple comparison test (***) ($P < 0.001$). Within the groups treated with different concentrations of Se-Met, Na₂SeO₃ or Se-Car, bars sharing a common letter are not significantly different ($P > 0.05$) (A). In the hepatocytes supplemented with Se as Se-Met, Na₂SeO₃ or Se-Car at the same dose, bars sharing a common letter are not significantly different ($P > 0.05$) (B).

3.2. Changes in intracellular GSH concentration

The effects of different forms and concentrations of Se on the intracellular concentration of GSH are shown in Fig. 2. In all Se-treated hepatocytes, intracellular concentrations of GSH were significantly lower than control ($P<0.001$). Increasing doses of Se-Met in the culture medium (1.5–5 $\mu\text{mol/L}$) did not lead to further significant reduction of intracellular GSH concentrations. In contrast, intracellular concentrations of GSH in the hepatocytes incubated with increased levels of Na_2SeO_3 or Se-Car were decreased in a dose-dependent manner.

3.3. Effect of Se supplementation on GPx1 mRNA level

The levels of GPx1 mRNA measured by quantitative RT-PCR are shown in Fig. 3. When compared with control, significant increases of GPx1 mRNA were observed in all groups treated with Se-Met, Na_2SeO_3 or Se-Car ($P<0.001$) (Fig. 3A). The maximal increases of GPx1 mRNA were observed in the hepatocytes incubated with Se-Met at a dose of 3 $\mu\text{mol/L}$ (about 6.54-fold vs. control), Na_2SeO_3 at a dose

of 1.5 $\mu\text{mol/L}$ (about 6.08-fold vs. control) or Se-Car at a dose of 2 $\mu\text{mol/L}$ (about 5.72-fold vs. control) (Fig. 3A). After reaching a maximal level, increased Se supplementation led to a reduction in GPx1 mRNA (Fig. 3A). The degree of reduction of GPx1 mRNA was relatively mild in the groups treated with Se-Met at doses of 4 and 5 $\mu\text{mol/L}$ when compared with the groups treated with Na_2SeO_3 and Se-Car at equivalent doses (Fig. 3B). Na_2SeO_3 had a relatively greater effect on GPx1 mRNA than Se-Met and Se-Car when supplemented at a level equivalent to a dose of Se of 1.5 $\mu\text{mol/L}$ ($P<0.05$) (Fig. 3B). Se-Met had greater effects on GPx1 mRNA than did Na_2SeO_3 and Se-Car at doses of 3, 4 and 5 $\mu\text{mol/L}$ ($P<0.001$) (Fig. 3B). Furthermore, 3 $\mu\text{mol/L}$ of Se from Se-Met resulted in the highest level of GPx1 mRNA of all the groups treated with Se (Fig. 3B).

3.4. Effect of Se supplementation on GPx1 activity

The effects of different forms and concentrations of Se on the activity of GPx1 are shown in Fig. 4. Compared with control, the supplementation of Se-Met, Na_2SeO_3 or Se-Car




Fig. 4. Effects of different forms and concentrations of Se on GPx1 activity in bovine hepatocytes. The primary cultured bovine hepatocyte monolayers were treated with 0 (control), 0.5, 1, 1.5, 2, 3, 4 or 5 $\mu\text{mol/L}$ of Se as Se-Met, Na_2SeO_3 or Se-Car for 24 h. The activity of GPx1 in hepatocyte cytosol was measured using a spectrophotometric method and expressed as units per gram of protein. Bars represent mean \pm S.D. of triplicate cultures. Bars with "****" are statistically significantly different from control by one-way ANOVA followed by Tukey's multiple comparison test ($***P<0.001$). Within the groups treated with different levels of Se-Met, Na_2SeO_3 or Se-Car, bars sharing a common letter are not significantly different ($P>0.05$) (A). In the hepatocytes supplemented with the same Se level, bars sharing a common letter are not significantly different ($P>0.05$) (B).

led to significant increase in GPx1 activity over the whole dose range ($P < 0.001$) (Fig. 4A). In groups treated with Se-Met, there was a significant dose-dependent increase in GPx1 activity (vs. control), up to a maximum at a dose of 3 $\mu\text{mol/L}$ (Fig. 4A). For Na_2SeO_3 supplementation, 1.5 $\mu\text{mol/L}$ was the most effective dose, and a decreasing trend in GPx1 activity was observed at doses of 1.5–5 $\mu\text{mol/L}$, in a dose-dependent manner (Fig. 4A). In the hepatocytes treated with Se-Car, the maximal increase of GPx1 activity was observed at a dose of 2 $\mu\text{mol/L}$, and higher Se-Car supplementation led to a reduction in GPx1 activity (Fig. 4A). At 1.5 $\mu\text{mol/L}$ of added Se, Na_2SeO_3 had a relatively greater effect on the activity of GPx1 than Se-Met and Se-Car ($P < 0.05$) (Fig. 4B). However, Se-Met had greater effects on the activity of GPx1 than Na_2SeO_3 and Se-Car at doses of 3, 4 and 5 $\mu\text{mol/L}$ ($P < 0.001$) (Fig. 4B). Furthermore, 3 $\mu\text{mol/L}$ of Se from Se-Met resulted in the highest level of activity of GPx1 (638.71 U/g protein) of all groups treated with Se (Fig. 4B). The degree of decrease in GPx1 activity was relatively mild in the hepatocytes incubated with Se-Met at doses of 4 and 5 $\mu\text{mol/L}$ when compared with that in the groups treated with Na_2SeO_3 and Se-Car at equivalent doses (Fig. 4B). The activity of GPx1 showed a similar pattern to that of the level of GPx1 mRNA.

4. Discussion

Several chemical forms of Se have been observed in the environment and in diets. Se occurs in the +6 oxidation state as selenate, the +4 oxidation state as selenite, the 0 oxidation state as elemental Se, the -1 oxidation state as selenocystine and the -2 oxidation state as Se-Met. Not all of these forms are metabolized in the same way [33]. The inorganic forms of Se (e.g., sodium selenite) are reduced to the selenide state using reducing equivalents from GSH and NADPH. In contrast, the organic forms of Se (e.g., Se-Met) release Se in the selenide state as a result of catabolism. Se-Met is activated initially by adenosylation and then demethylated and converted to Sec via selenohomocysteine and selenocystathionine in an analogous way to Met, and the Sec formed is degraded further in the liver to serine and selenide [5]. Furthermore, Se-Met can be incorporated nonspecifically into proteins in place of Met. Selenide plays an intermediary role in Se metabolism and is either used for selenoprotein synthesis or methylated to dimethyl selenide and trimethylselenonium ions and exhaled or excreted [5]. Because the metabolic pathways in the body are different for the inorganic and organic forms of Se [33], the absorption and bioavailability may vary. Previous data have shown that selenate has lower biological activity and absorption than selenite [34]. Organic forms of Se possess higher bioavailability [21,22] and lower toxicity compared with inorganic forms of Se [19,20]. The results of this study supported these findings.

LDH is a stable cytosolic enzyme, which becomes extracellular when the cell membrane is damaged. Therefore, the integrity and viability of the cell membrane, and the level of cytotoxicity, can be analyzed by determining the activity of LDH in the culture supernatant. Se is an essential trace element, but it can be toxic at a level not much higher than the beneficial dose. Earlier research has shown that incubation of rat hepatocytes for 2 h with Se as Na_2SeO_3 in concentrations higher than 6.3 $\mu\text{mol/L}$ led to a decrease in cell viability compared with control cells [35]. However, Park and Whanger [36] reported that isolated rat hepatocytes exposed to 200 $\mu\text{mol/L}$ of Se as Na_2SeO_3 for 1 h only showed a 6% increase of LDH activity in the culture medium compared with controls. In the present study, concentrations of Se, supplied as Na_2SeO_3 , of greater than 2 $\mu\text{mol/L}$ led to a significant dose-dependent increase in the release of LDH, when compared with controls (Fig. 1). Similar results were obtained in hepatocytes treated with Se-Car when the dose of Se was greater than 3 $\mu\text{mol/L}$ (Fig. 1). A major reason for this difference between experiments may be the duration of incubation. The toxicity of Se is very strongly related to the chemical form. Se from organic forms is less toxic than that from inorganic forms [18–20], and selenite (Se^{4+}) is more toxic than selenate (Se^{6+}) in vitro [37,38]. In this study, treatment with Se-Met (0.5–5 $\mu\text{mol/L}$), Na_2SeO_3 (0.5–1 $\mu\text{mol/L}$) or Se-Car (0.5 $\mu\text{mol/L}$) for 24 h caused a significant decrease in leakage of LDH compared with controls (Fig. 1), which suggests that supplementation of the culture medium with Se enhanced the membrane integrity and antioxidant ability of the cells. The application of 5 $\mu\text{mol/L}$ Se-Met, <2 $\mu\text{mol/L}$ Na_2SeO_3 and <3 $\mu\text{mol/L}$ Se-Car did not result in cytotoxicity. Müller and Pallauf [39] also reported that Se as Na_2SeO_3 at a dose of 100 ng/ml (~1.27 $\mu\text{mol/L}$) had no cytotoxicity for primary cultured rabbit hepatocytes. The release of LDH induced by Na_2SeO_3 at doses of 3, 4 or 5 $\mu\text{mol/L}$ was significantly higher than that induced by Se-Car at equivalent doses (Fig. 1). These results support the hypothesis that Se-Met is less toxic than Na_2SeO_3 and Se-Car and that Se-Car is slightly less toxic than Na_2SeO_3 .

A ubiquitous thiol-containing tripeptide, GSH is one of the most potent cellular antioxidants and is mainly associated with the thiol group of the cysteine residue [40]. It acts as a cosubstrate of GPx and plays an important role in the scavenging of ROS, free radicals and reactive metabolites that may be generated under various conditions. In addition, GSH is an indicator of oxidative stress: its level is often increased in tissues or cells as an adaptive response in increased oxidative stress. It has been reported that deficiency of Se in rats leads to a decrease in hepatic GPx1 activity, in conjunction with an increase in the activity of hepatic glutathione *S*-transferase, and affects GSH metabolism by increasing its synthesis and release in the liver with a concomitant increase of plasma GSH concentration [41]. In the present study, incubation of bovine hepatocytes for 24 h with Se supplied as Se-Met, Na_2SeO_3 or Se-Car at doses of 0.5–5 $\mu\text{mol/L}$ reduced intracellular GSH concentrations

compared with controls (Fig. 2). This observation may be attributable to the fact that Se supplementation of the culture medium enhanced the activity of antioxidant enzymes in the hepatocytes, and thus, oxidative stress was reduced. Previous research has indicated that increasing the Na₂SeO₃ concentration in the culture medium enhanced the activity of GPx1 and decreased the level of intracellular GSH in rabbit hepatocytes [39].

Previous research has also shown that the mRNA level and activity of GPx1 in Se-deficient rat livers were reduced to 6% and 1% of the Se-supplemented levels, respectively [32]. Supplementation with Se at 0.2 mg/kg of the diet was required to support the full expression of GPx1 in pig liver [26]. The maximal GPx1 activity in livers of both pregnant and lactating rats was obtained at 0.075 and 0.1 mg Se/kg of diet, respectively [27]. Using in vitro experiments, the activity of GPx1 was markedly enhanced by increasing the Se concentration in the culture medium of primary cultured rabbit hepatocytes [39], but we speculate that the GPx1 activity did not reach a plateau at 100 ng/ml (about 1.27 μmol/L) of Na₂SeO₃ supplementation. In the present study, the largest increases in the level and activity of GPx1 mRNA were observed in the primary cultured hepatocytes treated with Se-Met at a dose of 3 μmol/L, Na₂SeO₃ at a dose of 1.5 μmol/L or Se-Car at a dose of 2 μmol/L (Figs. 3A and 4A). However, recent results of the study by Romanowska et al. [42] indicated that the mRNA level and activity of GPx1 in HPL1D cells exhibited plateaus at 10 nM Na₂SeO₃ supplementation, with no further change at higher concentrations. Ebert et al. [43] reported that GPx1 activity was enhanced 1.8-fold in bone marrow stromal cells that were cultured in the presence of 100 nM Se supplied as Na₂SeO₃. Incubation of human umbilical vein endothelial cells with 1 μmol/L of Se supplied as Se-Met for 24 h led to a 65% increase in GPx1 activity but did not result in a significant change in the level of GPx1 mRNA [44]. In this study, after reaching the maximal effect, higher Se supplementation led to dose-dependent decreasing trends of the mRNA level and activity of GPx1 in the hepatocytes treated with Se-Met, Na₂SeO₃ or Se-Car (Figs. 3A and 4A). The decreasing trend was mild in hepatocytes treated with Se-Met, moderate in hepatocytes treated with Se-Car and severe in hepatocytes treated with Na₂SeO₃. Recent research has also shown that 1 μmol/L of Se supplied as Na₂SeO₃ significantly increased the activity of GPx1 in mouse hepatoma cells (about 2-fold), and GPx1 activity exhibited a decreasing trend at a dose of 2 μmol/L of Se supplied as Na₂SeO₃ [45]. Cytotoxicity was observed when Se-Car and Na₂SeO₃ were applied at doses over 3 and 2 μmol/L, respectively, while Se-Met, even at a dose of 5 μmol/L, did not show any cytotoxicity (Fig. 1). Thus, we attribute, at least in part, the discrepancy in the finding of a decreasing trend to the following facts: Se-Met is less toxic than Na₂SeO₃ and Se-Car, and Se-Car is slightly less toxic than Na₂SeO₃. Toxicity associated with Se leads to inhibition of cell growth [46], DNA fragmentation [47] and reduction of protein synthesis [38]. In addition, the effects of

the three forms of Se on GPx1 activity were not significantly different when supplemented at Se doses of 0.5 and 1 μmol/L in this study (Fig. 4B). The effect of Na₂SeO₃ at a dose of 1.5 μmol/L was significantly higher than that of Se-Met and Se-Car at equivalent doses (Fig. 4B) and reached plateaus. The greatest effects of Se-Met and Se-Car were observed at doses of 3 and 2 μmol/L, respectively. The level of GPx1 mRNA and the activity of GPx1 induced by 3 μmol/L of Se-Met were the highest of those seen in all the groups treated with Se. These results suggest that different forms and doses of Se have varied effects on the regulation of the mRNA level and activity of GPx1.

Taken together, our data demonstrate that supplementation of the culture medium with Se can enhance the mRNA level and activity of GPx1 in bovine hepatocytes significantly. The roles of different forms and concentrations of Se in the regulation of mRNA level and activity of GPx1 differ in cultured cells. The optimal concentrations that support the full expression of GPx1 in bovine hepatocytes of the various forms of Se are also different.

Acknowledgments

The authors are grateful to Dr. Huawei Zeng for his help in revision and Dr. Guangming Li for his technical help in determination of GPx1 mRNA.

The project was supported by the National Natural Science Foundation of China (Grant Nos. 30671547 and 30871892) and the Natural Science Foundation of Jiangsu Province (Grant No. BK2006144).

References

- [1] McKenzie RC, Rafferty TS, Beckett GJ. Selenium: an essential element for immune function. *Immunol Today* 1998;19:342–5.
- [2] Aréchiga CF, Ortíz O, Hansen PJ. Effect of prepartum injection of vitamin E and selenium on postpartum reproductive function of dairy cattle. *Theriogenology* 1994;41:1251–8.
- [3] Kaur P, Bansal MP. Effect of selenium-induced oxidative stress on the cell kinetics in testis and reproductive ability of male mice. *Nutrition* 2005;21:351–7.
- [4] Mahmoud KZ, Edens FW. Influence of organic selenium on hsp70 response of heat-stressed and enteropathogenic *Escherichia coli*-challenged broiler chickens (*Gallus gallus*). *Comp Biochem Physiol C Toxicol Pharmacol* 2005;141:69–75.
- [5] Schrauzer GN. Anticarcinogenic effects of selenium. *Cell Mol Life Sci* 2000;57:1864–73.
- [6] Whanger PD. Selenium and its relationship to cancer: an update. *Br J Nutr* 2004;91:11–28.
- [7] Combs Jr GF. Current evidence and research needs to support a health claim for selenium and cancer prevention. *J Nutr* 2005;135:343–7.
- [8] Tapiero H, Townsend DM, Tew KD. The antioxidant role of selenium and seleno-compounds. *Biomed Pharmacother* 2003;57:134–44.
- [9] Arthur JR, McKenzie RC, Beckett GJ. Selenium in the immune system. *J Nutr* 2003;133:1457S–9S.
- [10] The Research Group of Keshan Disease of Chinese Academy of Medical Sciences. Research on the relation between selenium and Keshan disease. *Acta Acad Med Sin* 1979;1:75–80.

- [11] Research Team of Environment and Endemic Diseases. Studies on low selenium belt in China and pathogeny of Keshan and Kaschin-Beck diseases. *Chin J Environ Sci* 1986;7:89–93.
- [12] Muth OH, Oldfield JE, Remmert LF, Schubert JR. Effects of selenium and vitamin E on white muscle disease. *Science* 1958;128:1090.
- [13] Patterson EL, Milstrey R, Stokstad EL. Effect of selenium in preventing exudative diathesis in chicks. *Proc Soc Exp Biol Med* 1957;95:617–20.
- [14] Rotruck JT, Pope AL, Ganther HE, Swanson AB, Hafeman DG, Hockstra WG. Selenium biochemical role as a component of glutathione peroxidase. *Science* 1973;179:588–90.
- [15] Arthur JR. The glutathione peroxidases. *Cell Mol Life Sci* 2000;57:1825–35.
- [16] Baker RD, Baker SS, LaRosa K, Whitney C, Newburger PE. Selenium regulation of glutathione peroxidase in human hepatoma cell line Hep3B. *Arch Biochem Biophys* 1993;304:53–7.
- [17] Moriarty PM, Reddy CC, Maquat LE. Selenium deficiency reduces the abundance of mRNA for Se-dependent glutathione peroxidase 1 by a UGA-dependent mechanism likely to be nonsense codon-mediated decay of cytoplasmic mRNA. *Mol Cell Biol* 1998;18:2932–9.
- [18] Mihajlović M. Selenium toxicity in domestic animals. *Glas Srp Akad Nauka (Med)* 1992;131–44.
- [19] Kim YY, Mahan DC. Comparative effects of high dietary levels of organic and inorganic selenium on selenium toxicity of growing–finishing pigs. *J Anim Sci* 2001;79:942–8.
- [20] Tiwary AK, Stegelmeier BL, Panter KE, James LF, Hall JO. Comparative toxicosis of sodium selenite and selenomethionine in lambs. *J Vet Diagn Invest* 2006;18:61–70.
- [21] Fairweather-Tait SJ. Bioavailability of selenium. *Eur J Clin Nutr* 1997;51:20S–3S.
- [22] Rayman MP. The use of high-selenium yeast to raise selenium status: how does it measure up. *Br J Nutr* 2004;92:557–73.
- [23] Sunde RA, Thompson BM, Palm MD, Weiss SL, Thompson KM, Evenson JK. Selenium regulation of selenium-dependent glutathione peroxidases in animals and transfected CHO cells. *Biomed Environ Sci* 1997;10:346–55.
- [24] Weiss Sachdev S, Sunde RA. Selenium regulation of transcript abundance and translational efficiency of glutathione peroxidase-1 and -4 in rat liver. *Biochem J* 2001;357:851–8.
- [25] Thomson CD. Assessment of requirements for selenium and adequacy of selenium status: a review. *Eur J Clin Nutr* 2004;58:391–402.
- [26] Lei XG, Dann HM, Ross DA, Cheng WH, Combs Jr GF, Roneker KR. Dietary selenium supplementation is required to support full expression of three selenium-dependent glutathione peroxidases in various tissues of weanling pigs. *J Nutr* 1998;128:130–5.
- [27] Sunde RA, Evenson JK, Thompson KM, Sachdev SW. Dietary selenium requirements based on glutathione peroxidase-1 activity and mRNA levels and other Se-dependent parameters are not increased by pregnancy and lactation in rats. *J Nutr* 2005;135:2144–50.
- [28] Seglen PO. Preparation of isolated rat liver cells. *Methods Cell Biol* 1976;13:29–83.
- [29] Donkin SS, Armentano LE. Preparation of extended in vitro cultures of bovine hepatocytes that are hormonally responsive. *J Anim Sci* 1993;71:2218–27.
- [30] Dringen R, Kussmaul L, Hamprecht B. Detoxification of exogenous hydrogen peroxide and organic hydroperoxides by cultured astroglial cells assessed by microtiter plate assay. *Brain Res Protoc* 1998;2:223–8.
- [31] Rahman I, Kode A, Biswas SK. Assay for quantitative determination of glutathione and glutathione disulfide levels using enzymatic recycling method. *Nat Protoc* 2006;1:3159–65.
- [32] Lei XG, Evenson JK, Thompson KM, Sunde RA. Glutathione peroxidase and phospholipid hydroperoxide glutathione peroxidase are differentially regulated in rats by dietary selenium. *J Nutr* 1995;125:1438–46.
- [33] Zeng H, Combs Jr GF. Selenium as an anticancer nutrient: roles in cell proliferation and tumor cell invasion. *J Nutr Biochem* 2008;19:1–7.
- [34] Vendeland SC, Deagen JT, Butler JA, Whanger PD. Uptake of selenite, selenomethionine and selenate by brush border membrane vesicles isolated from rat small intestine. *Biometals* 1994;7:305–12.
- [35] Bell RR, Nonavinakere VK, Soliman MR, Early II JL. Effect of in vitro treatment of rat hepatocytes with selenium, and/or cadmium on cell viability, glucose output, and cellular glutathione. *Toxicology* 1991;69:111–9.
- [36] Park YC, Whanger PD. Toxicity, metabolism and absorption of selenite by isolated rat hepatocytes. *Toxicology* 1995;100:151–62.
- [37] Danielsson BR, Danielson M, Khayat A, Wide M. Comparative embryotoxicity of selenite and selenate: uptake in murine embryonal and fetal tissues and effects on blastocysts and embryonic cells in vitro. *Toxicology* 1990;63:123–36.
- [38] Uezono Y, Toyohira Y, Yanagihara N, Wada A, Taniyama K. Inhibition by selenium compounds of catecholamine secretion due to inhibition of Ca²⁺ influx in cultured bovine adrenal chromaffin cells. *J Pharmacol Sci* 2006;101:223–9.
- [39] Müller AS, Pallauf J. Effect of increasing selenite concentrations, vitamin E supplementation and different fetal calf serum content on GPx1 activity in primary cultured rabbit hepatocytes. *J Trace Elem Med Biol* 2003;17:183–92.
- [40] Curello S, Ceconi C, Cargnoni A, Cornacchiari A, Ferrari R, Albertini A. Improved procedure for determining glutathione in plasma as an index of myocardial oxidative stress. *Clin Chem* 1987;33:1448–9.
- [41] Hill KE, Burk RF, Lane JM. Effect of selenium depletion and repletion on plasma glutathione and glutathione-dependent enzymes in the rat. *J Nutr* 1987;117:99–104.
- [42] Romanowska M, Kikawa KD, Fields JR, Maciag A, North SL, Shiao YH, Kasprzak KS, Anderson LM. Effects of selenium supplementation on expression of glutathione peroxidase isoforms in cultured human lung adenocarcinoma cell lines. *Lung Cancer* 2007;55:35–42.
- [43] Ebert R, Ulmer M, Zeck S, Meissner-Weigl J, Schneider D, Stopper H, Schupp N, Kassem M, Jakob F. Selenium supplementation restores the antioxidative capacity and prevents cell damage in bone marrow stromal cells in vitro. *Stem Cells* 2006;24:1226–35.
- [44] Jornot L, Junod AF. Differential regulation of glutathione peroxidase by selenomethionine and hyperoxia in endothelial cells. *Biochem J* 1995;306:581–7.
- [45] Keck AS, Finley JW. Aqueous extracts of selenium-fertilized broccoli increase selenoprotein activity and inhibit DNA single-strand breaks, but decrease the activity of quinone reductase in Hepa 1c1c7 cells. *Food Chem Toxicol* 2006;44:695–703.
- [46] Spyrou G, Björnstedt M, Skog S, Holmgren A. Selenite and selenate inhibit human lymphocyte growth via different mechanisms. *Cancer Res* 1996;56:4407–12.
- [47] Garberg P, Ståhl A, Warholm M, Högborg J. Studies of the role of DNA fragmentation in selenium toxicity. *Biochem Pharmacol* 1988;37:3401–6.